arXiv:1008.4869v2 [hep-ph] 30 Nov 2010

Higgs couplings in a model with triplets

Heather E. Logan* and Marc-André Roy
Ottawa-Carleton Institute for Physics,
Carleton University, Ottawa K1S 5B6 Canada

(Dated: November 30, 2010)

Abstract

We study the couplings of a CP-even neutral Higgs boson h in a model containing one scalar SU(2)$_L$ doublet, one real triplet, and one complex triplet with hypercharge 1. Because the two triplets contribute to the ρ parameter with opposite signs, the triplet vacuum expectation values can be sizable. We show that (i) the hWW and hZZ couplings can be larger than the corresponding values in the Standard Model, and (ii) the ratio of the WW and ZZ couplings of h can be different than the corresponding ratio in the Standard Model. Neither of these results can occur in models containing only Higgs doublets. We also compute the rates for $gg \to h \to WW$ and $gg \to h \to ZZ$ and find that, for reasonable parameter values and $M_h \sim$ 140–180 GeV, the hadron collider rate for $gg \to h \to WW$ (ZZ) can be up to 20% (5 times) larger than in the Standard Model. We discuss implications for Higgs coupling extraction at the LHC.

---

* logan@physics.carleton.ca

I. INTRODUCTION

The primary purpose of the CERN Large Hadron Collider (LHC) is to shed light on the dynamics of electroweak symmetry breaking. In the Standard Model (SM), this involves the discovery of the Higgs boson and the measurement of its mass-generating couplings to fermions and gauge bosons. Most extensions of the SM contain one or more Higgs-like scalars that play a similar role in mass generation; the characterization of such models also involves measurement of the couplings of these scalars to the SM fermions and gauge bosons.

The LHC will be able to measure Higgs signal rates in a variety of production and decay modes [1, 2], especially if the Higgs mass lies in the range 114 GeV $< M_h \lesssim$ 200 GeV, as suggested by direct searches [3] and electroweak precision data [4]. These signal rate measurements will provide access to the combinations of Higgs couplings that determine the production cross sections and decay branching ratios involved in each channel. By taking ratios of Higgs signal rates with either the production mode or the decay mode in common, ratios of Higgs couplings-squared can be found with no model-dependent assumptions [5, 6].

Unfolding the individual Higgs couplings from these rate measurements, however, is a major challenge. The main difficulty lies in the fact that the LHC will not provide an absolute Higgs production cross section measurement in any channel. Together with the possible presence of undetectable Higgs decay modes (e.g., $h \to gg$ or decays into light quarks), this leads to a degeneracy in the Higgs coupling fit corresponding to increasing all production couplings by a common factor while simultaneously decreasing the detectable branching fractions by the same factor. This forces the adoption of theoretical assumptions in Higgs coupling fits. Analyses of LHC Higgs coupling extraction prospects have dealt with this difficulty either by assuming that the total Higgs width is dominated by the known SM decay modes [5, 6], or by assuming that the couplings of the Higgs to WW and ZZ are bounded from above by their SM values [7–9] (the latter analyses make the simultaneous assumption that the ratio of the hWW and hZZ couplings is the same as in the SM). The latter approach allows the total width of the Higgs to be bounded from above by a rate measurement in a channel with Higgs production and decay via the hWW or hZZ coupling (e.g., weak boson fusion with $h \to WW$). These two assumptions about the hWW and hZZ couplings are valid in any Higgs sector that contains only $\mathrm{SU}(2)_L$ doublets and/or singlets. Larger $\mathrm{SU}(2)_L$ multiplets are usually considered unimportant because their contribution to the hWW and hZZ couplings are generally tightly constrained by the experimental limits on the ρ parameter.

In this paper we explore the possibility that these assumptions about the hWW and hZZ couplings are violated. To do so, we need a model with at least one Higgs doublet (required to generate fermion masses), together with at least one larger $\mathrm{SU}(2)_L$ representation (to give rise to group-theoretic coefficients in the hWW and hZZ couplings different from those of a doublet). Because the couplings of interest arise from quartic terms in the Lagrangian in which one scalar field is replaced by its vacuum expectation value (vev), we need the vev(s) of the larger $\mathrm{SU}(2)_L$ multiplet(s) to be non-negligible.

To this end we study a model containing one Higgs doublet and two triplets first introduced by Georgi and Machacek [10]. One triplet is real with hypercharge zero while the other is complex with hypercharge 1.[1] Because these two triplets contribute to the ρ parameter with opposite sign, their vevs need not be small so long as they are chosen such that their contributions to ρ cancel; this is achieved naturally by imposing a global "custodial" SU(2) symmetry [10–14]. (The required global symmetry is however broken by hypercharge, so it is no longer exact once radiative corrections are included [13].) The model contains three CP-even neutral scalars. We study one of these states, which we call h and parameterize as a general mixture of the three weak eigenstates. We find that the hWW and hZZ couplings can be significantly larger than their SM values, and that the ratio of these couplings can be significantly different from the ratio in the SM.

This paper is organized as follows. In the next section we summarize the model and present general expressions for the couplings of interest. In Sec. III we study the range of hWW and hZZ couplings allowed in the model, subject to experimental constraints that set a lower bound on the doublet vev. We also consider the impact on Higgs production and decay in two of the most promising early Higgs discovery channels at the LHC, $gg \to h \to WW$ and $gg \to h \to ZZ$. In particular, we find that the $gg \to h \to ZZ$ rate can be significantly enhanced compared to that in the SM, due to a suppression of the $h \to WW$ branching fraction in some parts of parameter space without a simultaneous suppression of $h \to ZZ$. In Sec. IV we discuss ways to experimentally determine whether the hZZ and hWW couplings have the same ratio as in the SM and whether these couplings are larger than their SM values. In Sec. V we summarize our conclusions. Some computational details are given in the appendices.

---

[1] We use the convention that the hypercharge of the SM Higgs doublet is 1/2.

II. THE MODEL

The Georgi-Machacek model [10] contains a complex SU(2)$_L$ scalar doublet Φ with hypercharge $Y = Q - T^3 = 1/2$, a real triplet Ξ with $Y = 0$, and a complex triplet X with $Y = 1$, with components and vevs denoted by

$$\Phi = \begin{pmatrix} \phi^+ \\ \frac{1}{\sqrt{2}}(v_\phi + \phi^{0,r} + i\phi^{0,i}) \end{pmatrix}, \qquad \Xi = \begin{pmatrix} \xi^+ \\ v_\xi + \xi^0 \\ \xi^- \end{pmatrix}, \qquad X = \begin{pmatrix} \chi^{++} \\ \chi^+ \\ \frac{1}{\sqrt{2}}(v_\chi + \chi^{0,r} + i\chi^{0,i}) \end{pmatrix}. \tag{1}$$

Starting from the canonically normalized covariant derivative terms in the scalar Lagrangian, we obtain the W and Z boson masses and their couplings to the CP-even neutral scalars as follows:

$$\begin{aligned} \mathcal{L} &\supset |\mathcal{D}_\mu\Phi|^2 + \frac{1}{2}|\mathcal{D}_\mu\Xi|^2 + |\mathcal{D}_\mu X|^2 \\ &\supset (v_\phi + \phi^{0,r})^2\left[\frac{g^2}{4}W^+_\mu W^{-\mu} + \frac{g^2+g'^2}{8}Z_\mu Z^\mu\right] \\ &\quad +(v_\xi + \xi^0)^2\left[4\,\frac{g^2}{4}W^+_\mu W^{-\mu}\right] \\ &\quad +(v_\chi + \chi^{0,r})^2\left[2\,\frac{g^2}{4}W^+_\mu W^{-\mu} + 4\,\frac{g^2+g'^2}{8}Z_\mu Z^\mu\right], \end{aligned} \tag{2}$$

where g and g' are the SU(2)$_L$ and U(1)$_Y$ gauge couplings of the SM, respectively. The covariant derivative and the SU(2) generators for the triplet representation are given in Appendix A. The squared masses of the W and Z bosons are then given by

$$M_W^2 = \frac{g^2}{4}(v_\phi^2 + 4v_\xi^2 + 2v_\chi^2) \equiv \frac{g^2}{4}v_{\rm SM}^2, \qquad M_Z^2 = \frac{g^2+g'^2}{4}(v_\phi^2 + 4v_\chi^2), \tag{3}$$

where we define $v_{\rm SM}^2 = v_\phi^2 + 4v_\xi^2 + 2v_\chi^2 \simeq (246\ {\rm GeV})^2$.

The different form of the triplet contributions to the W and Z masses results in a tree-level ρ parameter different from 1,

$$\rho \equiv \frac{M_W^2}{M_Z^2\cos^2\theta_W} = \frac{v_\phi^2 + 4v_\xi^2 + 2v_\chi^2}{v_\phi^2 + 4v_\chi^2} = 1 + \frac{4v_\xi^2 - 2v_\chi^2}{v_\phi^2 + 4v_\chi^2} \equiv 1 + \Delta\rho, \tag{4}$$

where we used the definition $\cos\theta_W = g/\sqrt{g^2+g'^2}$. The ρ parameter also receives radiative corrections in the SM from isospin violation, depending mostly on the top quark and SM

Higgs masses. After separating out the SM contributions, the constraint on new sources of SU(2) breaking is given at the 2σ level by [15]

$$\Delta\rho = 4^{+27}_{-7} \times 10^{-4} \qquad (2\sigma \text{ constraint}). \tag{5}$$

This leads to the requirement $v_\xi \simeq v_\chi/\sqrt{2}$ to high precision. This relation can be enforced by imposing an $\mathrm{SU}(2)_R$ global symmetry on the Higgs potential, resulting in a residual global "custodial" SU(2) symmetry—the diagonal subgroup of $\mathrm{SU}(2)_L\times \mathrm{SU}(2)_R$—after electroweak symmetry breaking [10–12]. The $\mathrm{SU}(2)_R$ global symmetry is broken by hypercharge (for the scalars, hypercharge corresponds to the T^3 generator of $\mathrm{SU}(2)_R$), so that $\mathrm{SU}(2)_R$-violating counterterms must be introduced at one-loop level [13]. Nevertheless, if the scale where $\mathrm{SU}(2)_R$ is a good symmetry is not too high, the approximate global symmetry protects the ρ parameter and has been used, e.g., to control the effects of triplets on electroweak precision observables in little Higgs models [16].

The imposition of custodial SU(2) on the Higgs potential results in Higgs mass eigenstates after electroweak symmetry breaking that lie in multiplets of the custodial SU(2) with common masses [10]. In particular, the physical states form two singlets H_1^0 and $H_1^{0\prime}$, a 3-plet $H_3^{+,0,-}$, and a 5-plet $H_5^{++,+,0,-,--}$ under custodial SU(2), with the CP-even neutral states given by [10]

$$H_1^0 = \phi^{0,r}, \qquad H_1^{0\prime} = \frac{1}{\sqrt{3}}(\xi^0 + \sqrt{2}\chi^{0,r}), \qquad H_5^0 = \frac{1}{\sqrt{3}}(\sqrt{2}\xi^0 - \chi^{0,r}). \tag{6}$$

If the custodial SU(2) is exact, the two singlets H_1^0 and $H_1^{0\prime}$ can mix, but H_5^0 must be a mass eigenstate.

In this framework, the couplings of the CP-even neutral states to W, Z, and fermion pairs are given by [10, 12]

$$\begin{aligned}
\bar g_{H_1^0WW} &= c_H, & \bar g_{H_1^0ZZ} &= c_H, & \bar g_{H_1^0ff} &= \frac{1}{c_H}, \\
\bar g_{H_1^{0\prime}WW} &= \frac{2\sqrt{2}}{\sqrt{3}}s_H, & \bar g_{H_1^{0\prime}ZZ} &= \frac{2\sqrt{2}}{\sqrt{3}}s_H, & \bar g_{H_1^{0\prime}ff} &= 0, \\
\bar g_{H_5^0WW} &= \frac{1}{\sqrt{3}}s_H, & \bar g_{H_5^0ZZ} &= -\frac{2}{\sqrt{3}}s_H, & \bar g_{H_5^0ff} &= 0,
\end{aligned} \tag{7}$$

where we define $\cos\theta_H \equiv c_H = v_\phi/v_{\mathrm{SM}}$, $s_H = \sqrt{1-c_H^2}$, and the barred couplings are normalized to the corresponding SM Higgs couplings, $\bar g_{hxx} \equiv g_{hxx}/g_{H_{\mathrm{SM}}xx}$.

In this paper we relax the assumption of custodial SU(2) in the Higgs potential. We still require that $v_\xi \simeq v_\chi/\sqrt{2}$ in order to obey the constraint on the ρ parameter, but we allow arbitrary mixing of the three CP-even neutral Higgs states. This is a fine-tuned situation, but it allows us to illustrate the full range of couplings allowed by experimental constraints. We will also present results for custodial SU(2)-preserving mixing. The most general gauge-invariant Higgs potential (without custodial SU(2)) is given in Appendix B, where we show that enough parameter freedom exists to allow arbitrary mixing even after the vevs are fixed as above.

The doublet and triplet vevs can be parameterized in terms of $v_{\rm SM}$, c_H, and $\Delta\rho$ according to

$$\begin{aligned} v_\phi^2 &= c_H^2 v_{\rm SM}^2, \\ v_\xi^2 &= \frac{1}{8}\left[s_H^2 + \frac{\Delta\rho}{1+\Delta\rho}\right] v_{\rm SM}^2 \simeq \frac{1}{8} s_H^2 v_{\rm SM}^2, \\ v_\chi^2 &= \frac{1}{4}\left[s_H^2 - \frac{\Delta\rho}{1+\Delta\rho}\right] v_{\rm SM}^2 \simeq \frac{1}{4} s_H^2 v_{\rm SM}^2. \end{aligned} \tag{8}$$

We consider two parameterizations of the mixing among the three neutral states. First, we parameterize the CP-even neutral Higgs state h of interest as

$$h = a\,\phi^{0,r} + b\,\xi^0 + c\,\chi^{0,r}, \tag{9}$$

where $a^2 + b^2 + c^2 = 1$.[2] The couplings of h are then given by

$$\bar g_{hWW} = \frac{a v_\phi + 4 b v_\xi + 2 c v_\chi}{v_{\rm SM}}, \qquad \bar g_{hZZ} = \frac{a v_\phi + 4 c v_\chi}{v_{\rm SM}}, \qquad \bar g_{hff} = \frac{a v_{\rm SM}}{v_\phi} = \frac{a}{c_H}. \tag{10}$$

In the limit $\Delta\rho \to 0$, the hWW and hZZ couplings can be written as

$$\bar g_{hWW} = a c_H + (\sqrt{2} b + c) s_H, \qquad \bar g_{hZZ} = a c_H + 2 c s_H. \tag{11}$$

As a second parameterization we can write

$$h = \cos\phi\,(\cos\theta\, H_1^0 + \sin\theta\, H_1^{0\prime}) + \sin\phi\, H_5^0, \tag{12}$$

where θ parameterizes the mixing between the two custodial SU(2) singlets and ϕ parameterizes the mixing between the custodial SU(2) singlets and the 5-plet. These mixing angles are related to a, b, and c above by

$$a = \cos\phi\cos\theta, \qquad b = \frac{1}{\sqrt{3}}\cos\phi\sin\theta + \sqrt{\frac{2}{3}}\sin\phi, \qquad c = \sqrt{\frac{2}{3}}\cos\phi\sin\theta - \frac{1}{\sqrt{3}}\sin\phi. \tag{13}$$

---

[2] In this notation, H_1^0, $H_1^{0\prime}$, and H_5^0 correspond to $(a,b,c) = (1,0,0)$, $(0,\sqrt{1/3},\sqrt{2/3})$, and $(0,\sqrt{2/3},-\sqrt{1/3})$, respectively.

In the limit $\Delta\rho = 0$ we have for the couplings,

$$\begin{aligned}
\bar{g}_{hWW} &= \cos\phi\cos\theta\, c_H + \cos\phi\sin\theta\frac{2\sqrt{2}}{\sqrt{3}}s_H + \sin\phi\frac{1}{\sqrt{3}}s_H, \\
\bar{g}_{hZZ} &= \cos\phi\cos\theta\, c_H + \cos\phi\sin\theta\frac{2\sqrt{2}}{\sqrt{3}}s_H - \sin\phi\frac{2}{\sqrt{3}}s_H, \\
\bar{g}_{hff} &= \cos\phi\cos\theta\frac{1}{c_H}.
\end{aligned} \tag{14}$$

Though less compact, this second parameterization will be more useful for our analysis. We note in particular that when $\sin\phi = 0$, h is a mixture of custodial SU(2) singlets only, and $\bar{g}_{hWW} = \bar{g}_{hZZ} \equiv \bar{g}_{hVV}$. For $\sin\phi = 0$ we have,

$$\bar{g}_{hVV} = \cos\theta\, c_H + \sin\theta\frac{2\sqrt{2}}{\sqrt{3}}s_H. \tag{15}$$

This coupling reaches a maximum value of

$$\bar{g}_{hVV}^{\max} = \sqrt{\frac{8-5c_H^2}{3}} \qquad \text{when} \qquad \sin\theta^{\max} = \sqrt{\frac{8s_H^2}{8-5c_H^2}}. \tag{16}$$

III. NUMERICAL RESULTS

Imposing $\Delta\rho = 0$, the largest hWW and hZZ couplings occur when $v_\phi = 0$ and $v_\chi = \sqrt{2}v_\xi = v_{\rm SM}/2$. This yields coupling maxima of $\bar{g}_{hWW} = \sqrt{3}$ for $h = \sqrt{\frac{2}{3}}\xi^0 + \frac{1}{\sqrt{3}}\chi^{0,r}$, for which $\bar{g}_{hZZ} = 2/\sqrt{3}$; and $\bar{g}_{hZZ} = 2$ for $h = \chi^{0,r}$, for which $\bar{g}_{hWW} = 1$. This situation is clearly unrealistic, however, because a nonzero value of v_ϕ is needed to generate fermion masses. This leads to a lower bound on c_H (equivalently v_ϕ) from processes that constrain the top quark Yukawa coupling in multi-Higgs-doublet models. In particular, the $\bar{b}b$ fraction in hadronic Z decays, R_b, was studied in the Georgi-Machacek model in Ref. [17]. For a custodial SU(2)-symmetric spectrum, $H_3^\pm$ contributes to R_b at one-loop level and leads to the constraint $c_H \geq 0.9$ (0.6) for $M_3 = 100$ (1000) GeV. If mixing between $H_3^\pm$ and $H_5^\pm$ is allowed, new contributions arise that can have either sign. In what follows we plot results for $c_H = 0.9$ and 0.95 as reasonable values. We also show the h couplings to WW and ZZ with $c_H = 0.6$ as an extreme case.

A. Higgs couplings

When plotted against each other for a fixed value of c_H, the expressions for $\bar{g}_{hWW}$ and $\bar{g}_{hZZ}$ given in Eqs. (11) and (14) describe a filled ellipse centered at the origin as shown in



FIG. 1. Normalized Higgs couplings $\bar{g}_{hZZ}$ versus $\bar{g}_{hWW}$ for a mixed state h in the Georgi-Machacek model, for $c_H = 0.6$, 0.9, and 0.95 (left to right). We sampled both signs for a, b, and c in Eq. (11). Points corresponding to a mixture of H_1^0 and $H_1^{0\prime}$ lie along the diagonal $\bar{g}_{hWW} = \bar{g}_{hZZ}$. H_5^0 is indicated by the crossed circle on the lower right boundary of the ellipse.

Fig. 1. The ellipse is bounded by a curve corresponding to $\theta = \theta^{\max}$ as defined in Eq. (16) and given parametrically by

$$\bar{g}_{hWW} = \cos\phi\sqrt{\frac{8-5c_H^2}{3}} + \sin\phi\frac{s_H}{\sqrt{3}},$$

$$\bar{g}_{hZZ} = \cos\phi\sqrt{\frac{8-5c_H^2}{3}} - \sin\phi\frac{2s_H}{\sqrt{3}}. \tag{17}$$

The bounding ellipse can be conveniently expressed by defining $g_\pm \equiv (\bar{g}_{hZZ} \pm \bar{g}_{hWW})/\sqrt{2}$; we then obtain

$$1 = \frac{3}{2(8-5c_H^2)}g_+^2 - \frac{1}{(8-5c_H^2)}g_+g_- + \frac{(11-7c_H^2)}{2s_H^2(8-5c_H^2)}g_-^2. \tag{18}$$

Note that this ellipse is not oriented along the diagonal $\bar{g}_{hWW} = \bar{g}_{hZZ}$, but instead has its semimajor axis tilted at a small angle φ above the diagonal, where

$$\tan 2\varphi = \frac{s_H^2}{2(1+s_H^2)}. \tag{19}$$

In the custodial SU(2)-symmetric case, H_5^0 must be a mass eigenstate but H_1^0 and $H_1^{0\prime}$ can mix. Points corresponding to $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$ fall along the diagonal, $\bar{g}_{hWW} = \bar{g}_{hZZ}$. The normalized couplings of H_5^0 are given in Table I. In particular, $\bar{g}_{H_5^0ZZ} = -2\bar{g}_{H_5^0WW}$. The $h = H_5^0$ point falls on the boundary of the ellipses in Fig. 1, marked with a crossed circle.

c_H	$\bar g_{H_5^0 WW}$	$\bar g_{H_5^0 ZZ}$
0.6	0.462	-0.924
0.9	0.252	-0.503
0.95	0.180	-0.361

TABLE I. Normalized couplings of H_5^0 to WW and ZZ for various values of c_H.

B. Higgs production and decay rates

We now compute the rates for $gg \to h \to WW$ and $gg \to h \to ZZ$, normalized to their SM values. In the SM, these processes constitute the most promising discovery modes for an intermediate-mass Higgs. Our calculation is valid for any hadron collider, because dependence on the initial-state parton densities and collider center-of-mass energy cancel in the ratio with the SM rate. We assume throughout that the narrow-width approximation can be applied. For $VV = WW$ or ZZ we have,

$$\sigma(gg \to h \to VV)/\sigma_{\rm SM} \equiv \frac{\sigma(gg \to h \to VV)}{\sigma_{\rm SM}(gg \to H \to VV)} = \frac{\sigma(gg \to h)}{\sigma_{\rm SM}(gg \to H)} \frac{{\rm BR}(h \to VV)}{{\rm BR}_{\rm SM}(H \to VV)}. \tag{20}$$

The $gg \to h$ production process proceeds through the coupling of h to a fermion loop and is dominated in the SM by the top quark contribution. Because all fermion couplings are scaled by the same factor $\bar g_{hff}$ in the Georgi-Machacek model, we have,[3]

$$\frac{\sigma(gg \to h)}{\sigma_{\rm SM}(gg \to H)} = \bar g_{hff}^2. \tag{21}$$

To compute the decay branching ratios in the Georgi-Machacek model, we start with the partial widths of the SM Higgs of the same mass, computed using the public FORTRAN code `HDECAY` version 3.531 [19]. The corresponding partial widths of h are computed as follows:

$$\Gamma(h \to WW) = \bar g_{hWW}^2 \Gamma_{\rm SM}(H \to WW), \qquad \Gamma(h \to ZZ) = \bar g_{hZZ}^2 \Gamma_{\rm SM}(H \to ZZ)$$
$$\Gamma(h \to f\bar f) = \bar g_{hff}^2 \Gamma_{\rm SM}(H \to f\bar f), \qquad \Gamma(h \to gg) = \bar g_{hff}^2 \Gamma_{\rm SM}(H \to gg), \tag{22}$$

---

[3] At two-loop order, there are electroweak contributions to the $gg \to h$ amplitude in which h couples to WW or ZZ inside the loop, which should instead be scaled by $\bar g_{hWW}$ or $\bar g_{hZZ}$, respectively. In the SM, these contributions result in a 5–6% positive correction to the cross section [18]. Ignoring their different coupling dependence thus results in an error of order $0.06\,\bar g_{hff}(\bar g_{hff} - \bar g_{hWW,hZZ})$ on the right-hand side of Eq. (21).



FIG. 2. The rate for $gg \to h \to WW$ normalized to its SM value plotted as a function of $\bar{g}^2_{hWW}$, for $M_h = 140$, 160 and 180 GeV (left to right), with $c_H = 0.9$. The solid line shows points for which $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$ and the crossed circle near the origin indicates the point corresponding to H_5^0. The dashed horizontal line shows the Tevatron upper limit on the rate for $gg \to h \to WW$ for the corresponding Higgs mass from Ref. [20].

where $f\bar{f}$ refers to any fermion pair. The remaining partial widths, $\Gamma(h \to \gamma\gamma)$ and $\Gamma(h \to \gamma Z)$, constitute a very small portion of the h total width; for simplicity we ignore the corrections to these modes and instead take them equal to the corresponding SM Higgs widths.[4] The branching ratios of h are then given by taking the ratio of the relevant partial width to the h total width.

We plot $\sigma(gg \to h \to WW)/\sigma_{\rm SM}$ in Figs. 2–5 and $\sigma(gg \to h \to ZZ)/\sigma_{\rm SM}$ in Figs. 6–9. We show results for $M_h = 140$, 160, and 180 GeV and $c_H = 0.9$ and 0.95. The points correspond to a general mixed state h, while the solid line shows the custodial SU(2)-preserving mixture $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$. For $h = H_5^0$, the $gg \to h$ cross section is zero because H_5^0 does not couple to fermions; this is shown in the plots by a crossed circle.

The rate for $gg \to h \to WW$ can be enhanced by up to about 20% (10%) compared to the corresponding SM rate for $c_H = 0.9$ (0.95), as shown in Table II. The enhancement is due to an interplay between the production and decay couplings. For $c_H = 0.9$ (0.95), the $gg \to h$ production cross section is about $1.23a^2$ ($1.11a^2$) times its SM value, for a defined in Eq. (9). The branching ratio for $h \to WW$ depends mainly on competition between the $h \to WW$ and $h \to \bar{b}b$ partial widths, especially for lower Higgs masses.

---

[4] The error introduced by this assumption in the h branching ratio calculation is at the level of $10^{-3}\,(1 - \bar{g}^2_{hff,hWW})$.



FIG. 3. As in Fig. 2 but for $c_H = 0.95$.



FIG. 4. The rate for $gg \to h \to WW$ normalized to its SM value plotted as a function of $\bar{g}^2_{hZZ}$, for $M_h = 140$, 160 and 180 GeV (left to right), with $c_H = 0.9$. The solid line shows points for which $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$ and the crossed circle at the lower edge of the allowed region indicates the point corresponding to H_5^0. The dashed horizontal line shows the Tevatron upper limit on the rate for $gg \to h \to WW$ for the corresponding Higgs mass from Ref. [20].



FIG. 5. As in Fig. 4 but for $c_H = 0.95$.



FIG. 6. The rate for $gg \to h \to ZZ$ normalized to its SM value plotted as a function of $\bar{g}_{hWW}^2$, for $M_h = 140$, 160, and 180 GeV (left to right), with $c_H = 0.9$. The solid line shows points for which $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$ and the crossed circle near the origin indicates the point corresponding to H_5^0. Points excluded by the Tevatron in Figs. 2 and 4 are not shown, leading to the gap in the solid line for $M_h = 160$ GeV.



FIG. 7. As in Fig. 6 but for $c_H = 0.95$.

For $c_H = 0.9$ and $M_h = 160$ GeV, the rate for $gg \to h \to WW$ can be enhanced enough to exceed the current limit on this cross section from the Tevatron [20].[5] Our calculation of the upper limit is given in Table III. Points excluded by the Tevatron are not shown in the plots of $gg \to h \to ZZ$ in Figs. 6–9.

The rate for $gg \to h \to ZZ$ can be enhanced much more dramatically, by more than a factor of 5 (3) for $M_h = 180$ (160) GeV and $c_H = 0.9$. Details are given in Table IV. This

---

[5] Note that the Tevatron analysis in Ref. [20] selected specifically for the $gg \to H \to WW$ process, and is not a combination of multiple SM Higgs search channels. The resulting cross section limits are thus directly applicable to the $gg \to h \to WW$ rates found here.



FIG. 8. The rate for $gg \to h \to ZZ$ normalized to its SM value plotted as a function of $\bar{g}_{hZZ}^2$, for $M_h = 140$, 160, and 180 GeV (left to right), with $c_H = 0.9$. The solid line shows points for which $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$ and the crossed circle at the lower edge of the allowed region indicates the point corresponding to H_5^0. Points excluded by the Tevatron in Figs. 2 and 4 are not shown, leading to the gap in the solid line for $M_h = 160$ GeV.



FIG. 9. As in Fig. 8 but for $c_H = 0.95$.

large enhancement occurs when the hWW coupling is near zero but the hZZ coupling is non-negligible, as can be seen by comparing Figs. 6–7 and Figs. 8–9. This suppresses the dominant Higgs partial width to WW, resulting in BR$(h \to ZZ)$ close to 1 for the heavier masses. Clearly, this can only happen if $\bar{g}_{hWW} \neq \bar{g}_{hZZ}$, so it requires mixing between H_5^0 and the custodial SU(2)-singlet states. This would lead to an early LHC discovery of such a Higgs in the "golden mode," $h \to ZZ \to 4$ leptons, while the $h \to WW$ signal (which would provide the first discovery in the SM) would be absent or very suppressed.

If $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$, the enhancement in the ZZ channel is limited to about 20%

		Maximum of $\sigma(gg \to h \to WW)/\sigma_{\rm SM}$	
c_H	M_h [GeV]	Overall	$h \sim H_1^0, H_1^{0\prime}$
0.9	140	1.1581	1.1352
	160	1.2085	1.2056
	180	1.2327	1.2296
0.95	140	1.0777	1.0656
	160	1.0971	1.0962
	180	1.1069	1.1061

TABLE II. Maximum values of $\sigma(gg \to h \to WW)/\sigma_{\rm SM}$ for $c_H = 0.9$ and 0.95 and $M_h = 140$, 160, and 180 GeV. The third column gives the maximum cross section enhancement obtainable when h is any linear combination of H_1^0, $H_1^{0\prime}$, and H_5^0, while the fourth column gives the maximum when h is a linear combination of only H_1^0 and $H_1^{0\prime}$.

	$M_h = 140$ GeV	160 GeV	180 GeV
$\sigma_{\rm SM}(gg \to H)$ [18]	0.680 pb	0.434 pb	0.279 pb
$\mathrm{BR}_{\rm SM}(H \to WW)$ [19]	0.4916	0.9048	0.9325
Tevatron upper limit on $\sigma(gg \to h \to WW)$ [20]	1.29 pb	0.47 pb	0.41 pb
Upper limit on $\sigma(gg \to h \to WW)/\sigma_{\rm SM}$	3.86	1.20	1.58

TABLE III. The $gg \to H$ cross section and $H \to WW$ branching fraction in the SM, Tevatron upper limit on the $gg \to h \to WW$ cross section, and resulting upper limit on the cross section normalized to its SM value, for $M_h = 140$, 160, and 180 GeV. The uncertainty on $\sigma_{\rm SM}(gg \to H)$ from the parton density functions is ±8–10% [18]; here we have taken the cross section central value for our limit.

(10%) for $c_H = 0.9$ (0.95). Note that when $h = \cos\theta H_1^0 + \sin\theta H_1^{0\prime}$, the enhancement in the WW and ZZ channels is identical, i.e., $\sigma(gg \to h \to WW)/\sigma_{\rm SM} = \sigma(gg \to h \to ZZ)/\sigma_{\rm SM}$. Such an enhancement would lead to a marginally earlier discovery of h in both channels at the LHC. For $c_H = 0.95$, however, the enhancement is about the same size as the current uncertainty in $\sigma_{\rm SM}(gg \to H)$ from the parton density functions [18].

		Maximum of $\sigma(gg \to h \to ZZ)/\sigma_{\rm SM}$	
c_H	M_h [GeV]	Overall	$h \sim H_1^0, H_1^{0\prime}$
0.9	140	1.5068	1.1356
	160	3.6616	1.1978
	180	5.7291	1.2297
0.95	140	1.2507	1.0659
	160	1.9716	1.0962
	180	3.0156	1.1061

TABLE IV. As in Table II but for $\sigma(gg \to h \to ZZ)/\sigma_{\rm SM}$.

IV. HIGGS COUPLING EXTRACTION AND THE TRIPLET NATURE OF h

In models containing only Higgs doublets and singlets, the hVV couplings satisfy

$$\bar g_{hWW} = \bar g_{hZZ} \equiv \bar g_{hVV}, \qquad |\bar g_{hVV}| \leq 1. \tag{23}$$

These relations are assumed to hold in the Higgs coupling fits of Refs. [7–9]; however, both of them are violated in the Georgi-Machacek model.

The relation $\bar g_{hWW} = \bar g_{hZZ}$ is easy to test by taking the ratio of any two Higgs signal rates with the same production mechanism and decays to WW and ZZ, respectively:

$$\frac{\sigma(X \to h \to WW)/\sigma_{\rm SM}(X \to h \to WW)}{\sigma(X \to h \to ZZ)/\sigma_{\rm SM}(X \to h \to ZZ)} = \frac{\bar g^2_{hWW}}{\bar g^2_{hZZ}}. \tag{24}$$

Determining whether $\bar g_{hVV}$ is greater than 1 is more complicated, and is not always possible at the LHC. The approach is as follows. One of the key ingredients in the Higgs coupling fits is the measurement of the rate for Higgs production in vector boson fusion (WW, $ZZ \to h$) followed by decay to WW. This rate can be written as

$$\sigma({\rm VBF} \to h \to WW) = \sigma({\rm VBF} \to h)\, {\rm BR}(h \to WW). \tag{25}$$

The usual Higgs coupling fit involves assuming that $\sigma({\rm VBF} \to h)$ is no larger than its SM value, i.e., $\sigma^{\rm max}({\rm VBF} \to h) = \sigma_{\rm SM}({\rm VBF} \to H)$. Together with the rate measurement, this sets a lower bound on ${\rm BR}(h \to WW)$:

$${\rm BR}^{\rm min}(h \to WW) = \frac{\sigma({\rm VBF} \to h \to WW)}{\sigma^{\rm max}({\rm VBF} \to h)}. \tag{26}$$

If the assumption that $\sigma(\mathrm{VBF} \to h) \leq \sigma_{\mathrm{SM}}(\mathrm{VBF} \to H)$ is false, the resulting value of $\mathrm{BR}^{\mathrm{min}}(h \to WW)$ can be greater than one, which is clearly unphysical. In particular, for $c_H = 0.9$ (0.95), we find that $\sigma(\mathrm{VBF} \to h)$ can be as much as 31% (16%) larger than in the SM. An unphysical $\mathrm{BR}^{\mathrm{min}}(h \to WW)$ would then be obtained using Eq. (26) if the true value of $\mathrm{BR}(h \to WW)$ is greater than 0.76 (0.86), which is true in the SM for M_H in the range 154–192 GeV (159–184 GeV).

This range can be extended by considering Higgs decays to other final states as follows. If the Higgs is detectable in the WW final state and some other final state (ZZ, $\tau\tau$, etc.) via a common production mechanism, we can write

$$\frac{\mathrm{BR}(h \to ZZ)}{\mathrm{BR}(h \to WW)} = \frac{\sigma(X \to h \to ZZ)}{\sigma(X \to h \to WW)}, \qquad \frac{\mathrm{BR}(h \to \tau\tau)}{\mathrm{BR}(h \to WW)} = \frac{\sigma(X \to h \to \tau\tau)}{\sigma(X \to h \to WW)}, \tag{27}$$

etc. The total branching fraction of the Higgs to detected modes is then given by

$$\mathrm{BR}(h \to \mathrm{detected}) = \mathrm{BR}(h \to WW) \left[1 + \frac{\mathrm{BR}(h \to ZZ)}{\mathrm{BR}(h \to WW)} + \frac{\mathrm{BR}(h \to \tau\tau)}{\mathrm{BR}(h \to WW)} + \cdots \right]. \tag{28}$$

If inserting $\mathrm{BR}^{\mathrm{min}}(h \to WW)$ from Eq. (26) on the right-hand side yields a value of $\mathrm{BR}(h \to \mathrm{detected})$ greater than 1, then we can conclude that the assumption that $\sigma(\mathrm{VBF} \to h) \leq \sigma_{\mathrm{SM}}(\mathrm{VBF} \to H)$ is false.

If the detectable branching fraction of the Higgs is sufficiently small, the technique discussed above will fail. In that case, detection of the triplet nature of h at the LHC would rely on the discovery of additional Higgs states. Search prospects at the LHC for single production of χ^{++} with decays to W^+W^+ were studied in Refs. [21, 22], and for pair production of $\chi^{++}\chi^{--}$ again with decays to like-sign W bosons in Ref. [23].[6] Single production of $H_5^+ \equiv (\chi^+ - \xi^+)/\sqrt{2}$ through W^+Z fusion with decays back to W^+Z was also studied in Refs. [22, 24].

We note that the technique discussed here can yield only a lower bound on $\sigma(\mathrm{VBF} \to h)$. Furthermore, removing the SM assumption that $\sigma(\mathrm{VBF} \to h) \leq \sigma_{\mathrm{SM}}(\mathrm{VBF} \to H)$ reopens the parameter degeneracy in the Higgs coupling fit discussed in the introduction. While ratios of Higgs couplings would still be measurable at the LHC, knowledge of absolute Higgs couplings would have to wait for direct model-independent measurements at an e^+e^- collider [25].

---

[6] Reference [23] also considered χ^{++} decays to like-sign charged leptons via lepton-number-violating Yukawa couplings that could be responsible for neutrino masses. For the triplet vevs that we consider here, $v_\chi = 54$ (38) GeV for $c_H = 0.9$ (0.95), such Yukawa couplings would be of order 10^{-11} and dilepton decays would be totally negligible.

V. CONCLUSIONS

We studied the couplings of a CP-even neutral Higgs state h in the Georgi-Machacek model with Higgs triplets. We found that the effect of the triplet components of h on its couplings to WW and ZZ can be significant, even for relatively small triplet vevs consistent with the experimental upper bounds on the top quark Yukawa coupling.

If h is a mixture of only custodial SU(2) singlets, the hWW and hZZ couplings can be enhanced by up to 15% (8%) for $c_H = 0.9$ (0.95), but the ratio of these couplings is the same as in the SM. If h is a generic mixture of the CP-even neutral states, the ratio of its WW and ZZ couplings can be very different than in the SM; in fact, one of these couplings can be zero while the other is finite. Such a suppression of the hWW coupling while the hZZ coupling is nonzero can lead to large enhancements in the rate for $gg \to h \to ZZ$, especially for $M_h \gtrsim 160$ GeV where the SM Higgs width is dominated by decays to WW.

While a ratio of hWW and hZZ couplings different than in the SM would be easy to measure by taking ratios of rates in these channels, an overall enhancement in the hVV coupling could only be detected at the LHC if it leads to a rate in vector boson fusion with decays to WW too large to be consistent with the SM assumption for this coupling.

ACKNOWLEDGMENTS

We thank K. Moats for helpful comments on the manuscript. This work was supported by the Natural Sciences and Engineering Research Council of Canada.

Appendix A: Covariant derivative and SU(2) generators for triplets

The gauge-covariant derivative in Eq. (2) is given by

$$\mathcal{D}_\mu = \partial_\mu + igT^a W^a_\mu + ig' Y B_\mu. \tag{A1}$$

The SU(2) generators for a doublet representation are given by $T^a = \sigma^a/2$, where σ^a are the 2×2 Pauli matrices. The SU(2) generators for a triplet representation are

$$T^1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \qquad T^2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \qquad T^3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \tag{A2}$$

Appendix B: Scalar potential and mass matrix

Reference [13] presented the most general gauge-invariant scalar potential for this model containing only terms with an even number of fields. While not completely general, this restricted potential captures most of the physics [11]. In our notation it is,

$$\begin{aligned}
V_{\rm even} = {} & \mu_1^2\Phi^\dagger\Phi + \mu_2^2 X^\dagger X + \mu_3^2\Xi^\dagger\Xi \\
& +\lambda_1(\Phi^\dagger\Phi)^2 + \lambda_2|X^T C X|^2 + \lambda_3(\Phi^\dagger T^a\Phi)(X^\dagger T^a X) \\
& +\lambda_4\left[(\Phi^\dagger T^a\Phi^c)(\Xi^\dagger T^a X) + \text{h.c.}\right] + \lambda_5(\Phi^\dagger\Phi)(X^\dagger X) + \lambda_6(\Phi^\dagger\Phi)(\Xi^\dagger\Xi) \\
& +\lambda_7(X^\dagger X)^2 + \lambda_8(\Xi^\dagger\Xi)^2 + \lambda_9|X^\dagger\Xi|^2 + \lambda_{10}(\Xi^\dagger\Xi)(X^\dagger X), \qquad \text{(B1)}
\end{aligned}$$

where λ_i are dimensionless and μ_i^2 have dimension mass squared. Here $\Phi^c = i\sigma_2\Phi^*$ and

$$C = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \qquad \text{(B2)}$$

is the conjugation operator for the triplet fields such that $X^c = CX^*$ transforms as a triplet of SU(2).

We add the most general gauge-invariant set of trilinear terms,

$$V_{\rm trilinear} = \kappa_1[(\Phi^\dagger T^a\Phi^c)X^a + \text{h.c.}] + \kappa_2(\Phi^\dagger T^a\Phi)\Xi^a + \kappa_3 X^{*a}(i\epsilon^{abc}X^b\Xi^c), \qquad \text{(B3)}$$

where κ_i have dimensions of mass and ϵ^{abc} is the totally antisymmetric tensor.

Minimizing the potential allows us to eliminate μ_i^2 in favor of the scalar vevs:

$$\begin{aligned}
\mu_1^2 &= -\kappa_1 v_\chi + \kappa_2\frac{v_\xi}{2} - \lambda_1 v_\phi^2 - \left(\frac{\lambda_3}{4} + \frac{\lambda_5}{2}\right)v_\chi^2 - \lambda_4 v_\chi v_\xi - \lambda_6 v_\xi^2 \\
\mu_2^2 &= -\kappa_1\frac{v_\phi^2}{2v_\chi} + \kappa_3 v_\xi - \left(\frac{\lambda_3}{4} + \frac{\lambda_5}{2}\right)v_\phi^2 - \lambda_4\frac{v_\phi^2 v_\xi}{2v_\chi} - \lambda_7 v_\chi^2 - \lambda_{10}v_\xi^2 \\
\mu_3^2 &= \kappa_2\frac{v_\phi^2}{8v_\xi} + \kappa_3\frac{v_\chi^2}{4v_\xi} - \lambda_4\frac{v_\phi^2 v_\chi}{4v_\xi} - \lambda_6\frac{v_\phi^2}{2} - 2\lambda_8 v_\xi^2 - \lambda_{10}\frac{v_\chi^2}{2}. \qquad \text{(B4)}
\end{aligned}$$

Applying these minimization conditions, the quadratic terms in the potential involving CP-even neutral fields are,

$$\begin{aligned}
V_{m^2\text{ terms}} = {} & (\phi^{0,r})^2\left[\lambda_1 v_\phi^2\right] \\
& +(\chi^{0,r})^2\left[-\kappa_1\frac{v_\phi^2}{4v_\chi} - \lambda_4\frac{v_\phi^2 v_\xi}{4v_\chi} + \lambda_7 v_\chi^2\right]
\end{aligned}$$

$$
\begin{aligned}
&+(\xi^0)^2\left[\kappa_2\frac{v_\phi^2}{8v_\xi}+\kappa_3\frac{v_\chi^2}{4v_\xi}-\lambda_4\frac{v_\phi^2 v_\chi}{4v_\xi}+4\lambda_8 v_\xi^2\right]\\
&+\phi^{0,r}\chi^{0,r}\left[\kappa_1 v_\phi+\left(\frac{\lambda_3}{2}+\lambda_5\right)v_\phi v_\chi+\lambda_4 v_\phi v_\xi\right]\\
&+\phi^{0,r}\xi^0\left[-\kappa_2\frac{v_\phi}{2}+\lambda_4 v_\phi v_\chi+2\lambda_6 v_\phi v_\xi\right]\\
&+\chi^{0,r}\xi^0\left[-\kappa_3 v_\chi+\frac{\lambda_4}{2}v_\phi^2+2\lambda_{10}v_\chi v_\xi\right]. \qquad \text{(B5)}
\end{aligned}
$$

Rewriting the fields in the H_1^0, $H_1^{0\prime}$, H_5^0 basis given in Eq. (6), using the notation of Eq. (8) for the vevs, and setting $\Delta\rho = 0$, we obtain

$$
\begin{aligned}
V_{m^2\ \mathrm{terms}} = v_{\mathrm{SM}}^2 \Bigg\{ &(H_1^0)^2\lambda_1 c_H^2\\
&+(H_1^{0\prime})^2\left[-\frac{(2\sqrt{2}\kappa_1-\kappa_2)}{v_{\mathrm{SM}}}\frac{c_H^2}{6\sqrt{2}s_H}-\frac{\kappa_3}{v_{\mathrm{SM}}}\frac{s_H}{4\sqrt{2}}+\lambda_4\frac{c_H^2}{3\sqrt{2}}+(\lambda_7+\lambda_8+\lambda_{10})\frac{s_H^2}{6}\right]\\
&+(H_5^0)^2\left[-\frac{(\kappa_1-\sqrt{2}\kappa_2)}{v_{\mathrm{SM}}}\frac{c_H^2}{6s_H}+\frac{\kappa_3}{v_{\mathrm{SM}}}\frac{s_H}{2\sqrt{2}}-\lambda_4\frac{3c_H^2}{4\sqrt{2}}+(\lambda_7+4\lambda_8-2\lambda_{10})\frac{s_H^2}{12}\right]\\
&+H_1^0H_1^{0\prime}\left[\frac{(2\sqrt{2}\kappa_1-\kappa_2)}{v_{\mathrm{SM}}}\frac{c_H}{2\sqrt{3}}+(\lambda_3+2\sqrt{2}\lambda_4+2\lambda_5+2\lambda_6)\frac{c_H s_H}{2\sqrt{6}}\right]\\
&+H_1^0H_5^0\left[-\frac{(\sqrt{2}\kappa_1+\kappa_2)}{v_{\mathrm{SM}}}\frac{c_H}{\sqrt{6}}+(-\lambda_3+\sqrt{2}\lambda_4-2\lambda_5+4\lambda_6)\frac{c_H s_H}{4\sqrt{3}}\right]\\
&+H_1^{0\prime}H_5^0\left[\frac{(\sqrt{2}\kappa_1+\kappa_2)}{v_{\mathrm{SM}}}\frac{c_H^2}{3s_H}+(-2\lambda_7+4\lambda_8+\lambda_{10})\frac{s_H^2}{6\sqrt{2}}\right]\Bigg\}. \qquad \text{(B6)}
\end{aligned}
$$

The last two terms, representing mixing between H_1^0 and H_5^0 and $H_1^{0\prime}$ and H_5^0, respectively, violate the custodial SU(2). Imposing the conditions given in Ref. [13] for the scalar potential to be invariant under a global $\mathrm{SU}(2)_R$ causes the λ_i contributions of these last two terms to vanish.[7] When $\kappa_2 = -\sqrt{2}\kappa_1$ the remaining contribution vanishes as well.

Even if the custodial SU(2)–violating couplings are small, the mixing between the singlets and H_5^0 can be large if their masses are degenerate enough.[8]

[1] G. Aad *et al.* [The ATLAS Collaboration], arXiv:0901.0512 [hep-ex].

[2] G. L. Bayatian *et al.* [CMS Collaboration], J. Phys. G **34**, 995 (2007).

---

[7] The relevant conditions are $\lambda_3 = \sqrt{2}\lambda_4$, $\lambda_5 = 2\lambda_6$, $2\lambda_7 = \lambda_9 + 2\lambda_{10}$, and $4\lambda_8 = \lambda_9 + \lambda_{10}$.

[8] The near-degenerate case was studied in a model with one doublet and one triplet in Ref. [26].

[3] R. Barate *et al.* [LEP Working Group for Higgs boson searches and ALEPH, DELPHI, L3, and OPAL Collaborations], Phys. Lett. B **565**, 61 (2003) [arXiv:hep-ex/0306033].

[4] J. Haller [The Gfitter group], arXiv:1006.0003 [hep-ph]; for updated results see `http://www.cern.ch/gfitter`.

[5] D. Zeppenfeld, R. Kinnunen, A. Nikitenko, and E. Richter-Was, in A. Djouadi *et al.*, arXiv:hep-ph/0002258; Phys. Rev. D **62**, 013009 (2000) [arXiv:hep-ph/0002036].

[6] A. Belyaev and L. Reina, JHEP **0208**, 041 (2002) [arXiv:hep-ph/0205270].

[7] M. Dührssen, ATL-PHYS-2003-003, available from `http://cdsweb.cern.ch`.

[8] M. Duhrssen, S. Heinemeyer, H. Logan, D. Rainwater, G. Weiglein and D. Zeppenfeld, Phys. Rev. D **70**, 113009 (2004) [arXiv:hep-ph/0406323].

[9] R. Lafaye, T. Plehn, M. Rauch, D. Zerwas and M. Duhrssen, JHEP **0908**, 009 (2009) [arXiv:0904.3866 [hep-ph]].

[10] H. Georgi and M. Machacek, Nucl. Phys. B **262**, 463 (1985).

[11] M. S. Chanowitz and M. Golden, Phys. Lett. B **165**, 105 (1985).

[12] J. F. Gunion, R. Vega and J. Wudka, Phys. Rev. D **42**, 1673 (1990).

[13] J. F. Gunion, R. Vega and J. Wudka, Phys. Rev. D **43**, 2322 (1991).

[14] J. F. Gunion, H. E. Haber, G. L. Kane and S. Dawson, *The Higgs Hunter's Guide* (Westview Press, Boulder, Colorado, 2000).

[15] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).

[16] S. Chang, JHEP **0312**, 057 (2003) [arXiv:hep-ph/0306034].

[17] H. E. Haber and H. E. Logan, Phys. Rev. D **62**, 015011 (2000) [arXiv:hep-ph/9909335].

[18] C. Anastasiou, R. Boughezal and F. Petriello, JHEP **0904**, 003 (2009) [arXiv:0811.3458 [hep-ph]].

[19] A. Djouadi, J. Kalinowski and M. Spira, Comput. Phys. Commun. **108**, 56 (1998) [arXiv:hep-ph/9704448], code available from `http://people.web.psi.ch/spira/hdecay/`.

[20] T. Aaltonen *et al.* [The CDF and D0 Collaborations], Phys. Rev. D **82**, 011102 (2010) [arXiv:1005.3216 [hep-ex]].

[21] G. Azuelos *et al.*, Eur. Phys. J. C **39S2**, 13 (2005) [arXiv:hep-ph/0402037].

[22] S. Godfrey and K. Moats, Phys. Rev. D **81**, 075026 (2010) [arXiv:1003.3033 [hep-ph]].

[23] T. Han, B. Mukhopadhyaya, Z. Si and K. Wang, Phys. Rev. D **76**, 075013 (2007) [arXiv:0706.0441 [hep-ph]].

[24] E. Asakawa, S. Kanemura and J. Kanzaki, Phys. Rev. D **75**, 075022 (2007) [arXiv:hep-ph/0612271].

[25] K. Desch [Higgs Working Group of the Extended ECFA/DESY Study], arXiv:hep-ph/0311092; T. L. Barklow, arXiv:hep-ph/0312268.

[26] A. G. Akeroyd and C. W. Chiang, Phys. Rev. D **81**, 115007 (2010) [arXiv:1003.3724 [hep-ph]].